



wwPDB X-ray Structure Validation Summary Report ⓘ

Jul 4, 2018 – 09:47 PM EDT

PDB ID : 1O7D
Title : The structure of the bovine lysosomal α -mannosidase suggests a novel mechanism for low pH activation
Authors : Heikinheimo, P.; Helland, R.; Leiros, H.S.; Leiros, I.; Karlsen, S.; Evjen, G.; Ravelli, R.; Schoehn, G.; Ruigrok, R.; Tollersrud, O.-K.; Mcsweeney, S.; Hough, E.
Deposited on : 2002-10-30
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : rb-20031172
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20031172


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2449 (2.70-2.70)
Clashscore	122126	2756 (2.70-2.70)
Ramachandran outliers	120053	2716 (2.70-2.70)
Sidechain outliers	120020	2716 (2.70-2.70)
RSRZ outliers	108989	2376 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	<div> <div>58%</div> <div>37%</div> <div>..</div> </div>
2	B	84	<div> <div>%</div> <div>48%</div> <div>38%</div> <div>5%</div> <div>10%</div> </div>
3	C	159	<div> <div>51%</div> <div>41%</div> <div>• 5%</div> </div>
4	D	282	<div> <div>48%</div> <div>46%</div> <div>..</div> </div>
5	E	126	<div> <div>48%</div> <div>34%</div> <div>• 13%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	NAG	C	1	X	-	-	-
6	NAG	D	11	X	-	-	-

2 Entry composition [i](#)

There are 12 unique types of molecules in this entry. The entry contains 7385 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lysosomal alpha-mannosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	292	Total	C	N	O	S	0	0	1
			2365	1520	408	426	11			

- Molecule 2 is a protein called Lysosomal alpha-mannosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	76	Total	C	N	O	S	0	0	1
			622	409	101	109	3			

- Molecule 3 is a protein called Lysosomal alpha-mannosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	151	Total	C	N	O	S	0	0	1
			1136	722	194	214	6			


- Molecule 4 is a protein called Lysosomal alpha-mannosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	270	Total	C	N	O	S	0	0	1
			2150	1356	397	392	5			

- Molecule 5 is a protein called Lysosomal alpha-mannosidase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	109	Total	C	N	O	S	0	0	1
			851	543	148	159	1			

- Molecule 6 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	C	1	Total	C	N	O	0	0
			14	8	1	5		
6	C	1	Total	C	N	O	0	0
			14	8	1	5		
6	D	1	Total	C	N	O	0	0
			14	8	1	5		
6	D	1	Total	C	N	O	0	0
			14	8	1	5		
6	D	1	Total	C	N	O	0	0
			14	8	1	5		
6	D	1	Total	C	N	O	0	0
			14	8	1	5		
6	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 7 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 8 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Zn	0	0
			1	1		

- Molecule 9 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	O	S	0	0
			5	4	1		
9	E	1	Total	O	S	0	0
			5	4	1		

- Molecule 10 is BETA-D-MANNOSE (three-letter code: BMA) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	C	1	Total	C	O	0	0
			11	6	5		

- Molecule 11 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	C	1	Total	C	O	0	0
			11	6	5		
11	C	1	Total	C	O	0	0
			11	6	5		
11	C	1	Total	C	O	0	0
			11	6	5		
11	C	1	Total	C	O	0	0
			11	6	5		
11	C	1	Total	C	O	0	0
			11	6	5		


- Molecule 12 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	14	Total	O	0	0
			14	14		
12	B	2	Total	O	0	0
			2	2		
12	C	3	Total	O	0	0
			3	3		
12	D	14	Total	O	0	0
			14	14		
12	E	6	Total	O	0	0
			6	6		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Lysosomal alpha-mannosidase



• Molecule 2: Lysosomal alpha-mannosidase




• Molecule 3: Lysosomal alpha-mannosidase




• Molecule 4: Lysosomal alpha-mannosidase





● Molecule 5: Lysosomal alpha-mannosidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	117.88Å 117.88Å 582.04Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.70 29.62 – 2.51	Depositor EDS
% Data completeness (in resolution range)	89.4 (30.00-2.70) 86.0 (29.62-2.51)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 2.51Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.257 , 0.289 0.252 , 0.249	Depositor DCC
R_{free} test set	3584 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	45.3	Xtriage
Anisotropy	0.085	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 45.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.37$, $\langle L^2 \rangle = 0.20$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	7385	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, BMA, NAG, SO4, TRS, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/2431	0.72	2/3302 (0.1%)
2	B	0.59	0/644	0.69	0/879
3	C	0.51	0/1163	0.72	0/1589
4	D	0.49	0/2198	0.74	0/2986
5	E	0.53	0/868	0.77	1/1186 (0.1%)
All	All	0.50	0/7304	0.73	3/9942 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	247	PRO	C-N-CD	-13.16	91.66	120.60
5	E	905	LEU	N-CA-C	-5.08	97.30	111.00
1	A	247	PRO	C-N-CA	5.05	143.23	122.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2365	0	2290	129	0
2	B	622	0	587	48	0
3	C	1136	0	1106	91	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	2150	0	2110	147	0
5	E	851	0	853	64	0
6	A	14	0	13	1	0
6	C	28	0	24	8	0
6	D	84	0	75	10	0
7	A	8	0	11	1	0
8	A	1	0	0	0	0
9	A	5	0	0	0	0
9	E	5	0	0	0	0
10	C	11	0	8	0	0
11	C	66	0	56	0	0
12	A	14	0	0	1	0
12	B	2	0	0	0	0
12	C	3	0	0	1	0
12	D	14	0	0	0	0
12	E	6	0	0	0	0
All	All	7385	0	7133	409	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 409 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:540:PRO:CA	5:E:887:THR:HG21	1.81	1.10
1:A:237:GLN:NE2	2:B:373:LYS:HB2	1.69	1.07
3:C:540:PRO:HA	5:E:887:THR:HG21	1.12	1.06
6:C:1:NAG:H81	4:D:839:ARG:H	1.20	1.04
2:B:403:ARG:HG3	2:B:403:ARG:HH11	1.14	1.02

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	290/298 (97%)	262 (90%)	25 (9%)	3 (1%)	17	41
2	B	74/84 (88%)	68 (92%)	5 (7%)	1 (1%)	12	31
3	C	149/159 (94%)	133 (89%)	15 (10%)	1 (1%)	24	50
4	D	266/282 (94%)	247 (93%)	16 (6%)	3 (1%)	16	38
5	E	105/126 (83%)	98 (93%)	6 (6%)	1 (1%)	17	41
All	All	884/949 (93%)	808 (91%)	67 (8%)	9 (1%)	17	41

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
5	E	920	GLN
1	A	77	TRP
2	B	421	PRO
4	D	874	GLY
1	A	260	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	251/260 (96%)	241 (96%)	10 (4%)	34	64
2	B	65/71 (92%)	59 (91%)	6 (9%)	10	23
3	C	125/139 (90%)	121 (97%)	4 (3%)	42	73
4	D	227/240 (95%)	218 (96%)	9 (4%)	34	64
5	E	93/111 (84%)	89 (96%)	4 (4%)	32	61
All	All	761/821 (93%)	728 (96%)	33 (4%)	32	61

5 of 33 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	409	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	536	LEU
5	E	907	ARG
2	B	410	GLN
3	C	472	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	511	GLN
4	D	628	ASN
4	D	779	ASN
2	B	410	GLN
3	C	445	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 1 is monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	A	1	1	14,14,15	0.70	0	17,19,21	0.80	0
7	TRS	A	2	8	7,7,7	1.89	3 (42%)	9,9,9	1.02	0
9	SO4	A	3001	-	4,4,4	0.62	0	6,6,6	0.26	0
6	NAG	C	1	3,6	14,14,15	0.60	0	17,19,21	1.28	1 (5%)
11	MAN	C	13	11,10	11,11,12	0.82	1 (9%)	15,15,17	0.90	0
11	MAN	C	14	11	11,11,12	1.66	2 (18%)	15,15,17	2.99	4 (26%)
11	MAN	C	16	11,10	11,11,12	0.46	0	15,15,17	0.87	1 (6%)
11	MAN	C	163	11	11,11,12	0.56	0	15,15,17	0.51	0
11	MAN	C	166	11	11,11,12	0.52	0	15,15,17	0.71	0
11	MAN	C	167	11	11,11,12	0.40	0	15,15,17	0.71	1 (6%)
6	NAG	C	2	10,6	14,14,15	0.56	0	17,19,21	0.86	1 (5%)
10	BMA	C	3	11,6	11,11,12	0.42	0	15,15,17	0.61	0
6	NAG	D	1	4,6	14,14,15	0.75	0	17,19,21	0.96	1 (5%)
6	NAG	D	11	6	14,14,15	0.82	0	17,19,21	1.18	2 (11%)
6	NAG	D	2	4,6	14,14,15	0.53	0	17,19,21	0.94	1 (5%)
6	NAG	D	21	6	14,14,15	0.58	0	17,19,21	0.75	0
6	NAG	D	3	4,6	14,14,15	0.74	0	17,19,21	1.08	2 (11%)
6	NAG	D	31	6	14,14,15	0.61	0	17,19,21	0.87	1 (5%)
9	SO4	E	3002	-	4,4,4	0.74	0	6,6,6	0.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	A	1	1	-	2/6/23/26	0/1/1/1
7	TRS	A	2	8	-	0/9/9/9	0/0/0/0
9	SO4	A	3001	-	-	0/0/0/0	0/0/0/0
6	NAG	C	1	3,6	1/1/5/7	0/6/23/26	0/1/1/1
11	MAN	C	13	11,10	-	0/2/19/22	0/1/1/1
11	MAN	C	14	11	-	0/2/19/22	0/1/1/1
11	MAN	C	16	11,10	-	0/2/19/22	0/1/1/1
11	MAN	C	163	11	-	0/2/19/22	0/1/1/1
11	MAN	C	166	11	-	0/2/19/22	0/1/1/1
11	MAN	C	167	11	-	0/2/19/22	0/1/1/1
6	NAG	C	2	10,6	-	0/6/23/26	0/1/1/1
10	BMA	C	3	11,6	-	0/2/19/22	0/1/1/1
6	NAG	D	1	4,6	-	0/6/23/26	0/1/1/1
6	NAG	D	11	6	1/1/5/7	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	D	2	4,6	-	0/6/23/26	0/1/1/1
6	NAG	D	21	6	-	0/6/23/26	0/1/1/1
6	NAG	D	3	4,6	-	0/6/23/26	0/1/1/1
6	NAG	D	31	6	-	0/6/23/26	0/1/1/1
9	SO4	E	3002	-	-	0/0/0/0	0/0/0/0

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	C	14	MAN	C1-C2	-3.22	1.44	1.52
7	A	2	TRS	C-N	-2.07	1.42	1.49
11	C	13	MAN	C2-C3	2.27	1.55	1.52
7	A	2	TRS	C3-C	2.71	1.59	1.52
7	A	2	TRS	C2-C	3.04	1.60	1.52

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	C	14	MAN	O3-C3-C2	-4.04	102.52	110.04
6	C	2	NAG	C2-N2-C7	-2.66	119.06	122.94
6	D	11	NAG	C2-N2-C7	-2.42	119.41	122.94
6	D	31	NAG	C2-N2-C7	-2.19	119.75	122.94
6	D	3	NAG	C2-N2-C7	-2.09	119.89	122.94

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	D	11	NAG	C1
6	C	1	NAG	C1

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	1	NAG	C8-C7-N2-C2
6	A	1	NAG	O7-C7-N2-C2

There are no ring outliers.

10 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1	NAG	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	2	TRS	1	0
6	C	1	NAG	5	0
6	C	2	NAG	4	0
6	D	1	NAG	3	0
6	D	11	NAG	3	0
6	D	2	NAG	4	0
6	D	21	NAG	2	0
6	D	3	NAG	1	0
6	D	31	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	292/298 (97%)	-0.59	0 100 100	22, 36, 54, 76	0
2	B	76/84 (90%)	-0.51	1 (1%) 77 78	18, 35, 70, 87	0
3	C	151/159 (94%)	-0.67	0 100 100	22, 36, 56, 95	0
4	D	270/282 (95%)	-0.54	1 (0%) 92 93	21, 38, 58, 94	0
5	E	109/126 (86%)	-0.66	0 100 100	16, 38, 59, 92	0
All	All	898/949 (94%)	-0.59	2 (0%) 94 96	16, 37, 58, 95	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	422	ALA	3.6
4	D	875	GLY	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
11	MAN	C	14	11/12	0.83	0.18	72,72,72,72	0
6	NAG	A	1	14/15	0.84	0.32	57,57,57,57	0
6	NAG	D	11	14/15	0.85	0.32	79,79,79,79	0
7	TRS	A	2	8/8	0.87	0.23	35,35,35,35	0
11	MAN	C	163	11/12	0.89	0.19	57,57,57,57	0
6	NAG	D	21	14/15	0.90	0.22	67,67,67,67	0
6	NAG	D	31	14/15	0.91	0.28	75,75,75,75	0
11	MAN	C	166	11/12	0.92	0.14	32,32,32,32	0
6	NAG	D	2	14/15	0.92	0.10	46,46,46,46	0
6	NAG	D	1	14/15	0.93	0.18	50,50,50,50	0
11	MAN	C	16	11/12	0.94	0.11	33,33,33,33	0
6	NAG	D	3	14/15	0.94	0.20	44,44,44,44	0
6	NAG	C	1	14/15	0.95	0.12	31,31,31,31	0
11	MAN	C	13	11/12	0.95	0.14	44,44,44,44	0
6	NAG	C	2	14/15	0.96	0.10	25,25,25,25	0
11	MAN	C	167	11/12	0.96	0.10	27,27,27,27	0
9	SO4	A	3001	5/5	0.96	0.20	36,36,36,36	0
10	BMA	C	3	11/12	0.97	0.10	33,33,33,33	0
9	SO4	E	3002	5/5	0.98	0.14	36,36,36,36	0
8	ZN	A	3	1/1	0.99	0.11	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.